

# Overview the Condition of Seawaters Around the Fish Landing Base in Sarang Indonesia

*by* Slamet Isworo

---

**Submission date:** 13-Mar-2020 12:13PM (UTC+0700)

**Submission ID:** 1274816560

**File name:** 6.\_Artikel\_Overview\_Seawaters.pdf (740.34K)

**Word count:** 8336

**Character count:** 42818

## Overview the Condition of Seawaters Around the Fish Landing Base in Sarang - Indonesia

Slamet Isworo<sup>(1\*)</sup> and Poerna Sri Oetari<sup>(2,3)</sup>

<sup>1</sup>Department of Health, Dian Nuswantoro University, Semarang, Indonesia.

<sup>2</sup>Mitra Adi Pranata Company, Environmental Impact Assessment (EIA) Consultants, Semarang, Indonesia.

<sup>3</sup>Graduate School of Environmental Science, Diponegoro University, Semarang, Indonesia.

\*Corresponding Author

### Abstract

**Background and Objective:** An Overview about the status of the waters of Sarang, it must be known from the beginning that the changes that occur due to the activities of fish landing bases so that it can be known and managed comprehensively. This study is a case study that produces an analysis that can be utilized by the local government in determining the development of its territory

**Materials and methods:** Tools and Materials: pH-meter, GPS, Thermometer, Secchi, Turbidimeter, Analytical Scale And Spectrophotometer, Eckman Grab. HNO<sub>3</sub>, Plankton Net, Dark Sample Bottle, Aquades, Formalin 4%,

Method. Analysis of sea air quality using AAS method and plankton, benthos and nekton analysis using Saprobity and Shannon Winner diversity index

**Results:** The results of water quality analysis found some indicators that exceed the quality standard limits of lead 0.9 ppm - 0.15 ppm, coliform  $4 \times 10^3$  -  $14 \times 10^3$  (MPN 71.00 mL), Ammonia = 519 ppm - 655 ppm, Phenol = 0.02 ppm - 0.22 ppm, H<sub>2</sub>S = 0.57 ppm - 0.74 ppm. Analysis of the biological component as follows Saprobik Index is 0.1, Tropic Saprobik Index is 0.24-0.49 and Nekton's Diversity Index is 1.74

**Conclusion:** Based on the category of water pollution levels around the Sarang fish landing base indicates that the ecological status of the category is moderately polluted.

**Keyword:** Plankton, Benthos, Nekton, Sarang of Seawaters, Quality of Seawaters, Diversity, Saprobity, AAS, Ecological status, Transect

### INTRODUCTION

Rembang regency is one of the Districts in the Republic of Indonesia, which has huge fishery and marine potential and has a long coastal area of ± 62.33 km. With the abundant potential of marine products, and supported by the presence of several residents who live along the coast with livelihoods as fishermen. Based on the statistical data of Rembang district stated that the total production of marine fishery in Rembang regency reach 65,523 tons. Given the second largest fishery resources in the province of Central Java, marine fisheries sector in Rembang waters is very potential in supporting the economy in Indonesia. Efforts of mining in supporting the distribution of fish catch by fishermen, fish landing facility

required, including fish auction place to accommodate and distribute the catch to the area around the province of Central Java. Fish landing activities Sarang located in Rembang regency has a land area of 2 hectares with an adequate dock.<sup>(1)</sup> Relation to the strategic potential of marine fishery products in Rembang region, it is necessary to develop based on regulation through Minister of Marine Affairs and Fisheries Decree no. 45 / KEPMEN-KP / 2014 on the Port Master Development Plan, stating that the status of Fishery Foundation is defined as Fish Landing Base from 2015 to 2034.<sup>(2)</sup> States that the development of Sarang fish landing base has been in accordance with the spatial plan of Rembang Regency, so it is necessary to study the compress environment in its development implementation, ecological condition status. Sarang waters need to know in advance so that the impact of fish landing development can be managed better<sup>(3)</sup>.

Pollution to marine waters will be more significant in the development and activity of fish landing base, this is because many organic and anorganic materials that can contaminate the sea due to fish landing activity at the port. pollutant materials from fisherman activities will be more accumulated at the port, thus causing disruption of ecological systems in coastal waters.<sup>(4)</sup>

According to Indicator of the quality of marine waters around the fish landing docks, it is important to consider the environmental parameters that must be managed and monitored as a standard quality standard set by the government<sup>(5)</sup> as follows:

1. **Water temperature** has a significant influence on the process of exchange or metabolism of living things. So that temperature is one important factor in regulating the process of life and the spread of organisms. The fish can grow well in 25-32 ° C.

2. **Total Suspended Solid (TSS)** a water sample is the amount of weight of a suspended material in a given water volume, with units of mg per liter.

Suspended solids consist of precipitated components, floating materials and suspended colloidal components. Suspended solids contain inorganic materials and organic materials. Therefore, TSS is an indicator of water soluble organic material. Respiratory aquatic organisms.<sup>(6)</sup>

3. **The degree of acidity (pH)** has a major influence on the life of aquatic organisms, so that the pH of a waters is often used as a guide to poor water quality. Aquatic organisms require water

conditions with a certain degree of acidity, water with a pH too high or too low can kill organisms, as well as changes. The organisms of marine or coastal ecosystems are relatively more stable and are within a narrow range, typically ranging from 7.7- 8.4.

**4. Phosphate** is one of the nutrients needed for the growth and metabolism of phytoplankton and marine organisms, and also determines the fertility of the waters. The elements of nitrogen and phosphorus present in the form of nitrate (NO<sub>3</sub>) and phosphate (PO<sub>4</sub>), under normal conditions at sea have a ratio of N / P is 15: 1. High levels of phosphate caused by the degradation of decomposed organic matter in marine sediments. Therefore, all the phosphates present in the underwater will rise to the surface, so that the surface layer becomes fertile resulting in enrichment (eutrophication) of nutrients in the ocean waters <sup>(7)</sup>.

**5. Nitrate.** Nutrients are needed in the growth of phytoplankton and other microorganisms as food sources and can also affect the fertility of marine waters. The eutrophication process in many waters is influenced by organic matter content.<sup>98</sup> <sup>(8,9)</sup> Nitrites are usually found in very small amounts of natural water, their levels are less than nitrate because nitrite is unstable when there is oxygen. Nitrite is a transitional form between ammonia and nitrite and between nitrate and oxygen gas commonly known as nitrification and denitrification in the biogeochemical cycle <sup>(10)</sup>.

**6. Salinity** has an important role and has a close relationship with the life of aquatic organisms including fish. The physiological salinity corresponds closely to the adjustment of fishing osmotic pressure. Salinity in waters farther from the mainland the higher the salinity so that the pH will become more alkaline. Salinity itself is the level of salinity or salt content in water. Salinity sea water has a salt content of 3-4 PSU equivalent to salinity 30-40 PSU (practical salinity units)<sup>(10)</sup>

**7. BOD (Biological Oxygen Demand)** BOD is one of the parameters that states the amount of oxygen needed by the organism in its metabolic processes. High BOD values indicate an increase in the amount of decomposed organic matter by <sup>36</sup> ng oxygen in water. The amount of BOD value can also be used to determine the presence of contamination in the water. So, more organic matter in the water so that the oxygen content is less soluble in it. Marine waters containing organic matter and suspended solids generally cause high BOD. <sup>46</sup>

**8. COD (Chemical Oxygen Demand)** COD represents the total amount of oxygen required to oxidize organic chemicals, but because of the high chloride content (Cl) in sea water so that the COD parameter is difficult to use as an indicator of marine pollution <sup>(11)</sup>

**9. Ammonia and Total Nitrogen.** Ammonia is one of the parameters of organic pollution in waters produced through the process of decomposition of organic matter (eutrophication) anaerobically by microbes. The high ammonia content in the waters will cause the water color to become turbid and produce odor. <sup>(11)</sup>

An ecosystem of biotic, abiotic, organic and inorganic factors interconnected to form a complex interaction relationship

pattern, as follows: high pH will cause the balance between ammonium and total ammonia in the water to be disturbed, the increase in pH will increase the total ammonia concentration is also very toxic to organisms. High ammonia concentrations on the surface of the water will cause fish deaths in the waters. The acidity of the water or its pH value greatly affects whether the amount of ammonia present will be toxic or not. The effect of pH on total ammonia toxicity is indicated by the state at low pH conditions that are toxic if the amount of ammonia is large, where <sup>8</sup> at high pH only with low amounts of ammonia will be toxic. High levels of ammonia can be used as an indication of contamination of organic materials derived from domestic waste and agricultural fertilizer runoff. The source of ammonia in the waters is the result of organic nitrogen breakdown in the form of dead water plants and aquatic biota. Fish can not tolerate too high concentrations of ammonia because it can interfere with the process of binding oxygen by the blood and can ultimately lead to death. In addition, ammonia also depends on the water temperature, <sup>(12)</sup>. The non-ionized ammonia form (NH<sub>3</sub>) is highly toxic while the ionized form (NH<sub>4</sub> + ion) is not toxic. These two forms are collectively called total ammonia <sup>(13)</sup>

**10. Sulfate.** Sulfate compound is found as a result of decomposition of organic matter. High sulfate levels in these waters are influenced by human activities carried through rivers to rivers and flow into coastal and marine waters <sup>(14)</sup>. These parameters can cause ecological imbalances, especially in marine waters around the Sarang fish landing port.

Plankton, Benthos and nekton are indicators of marine waters, therefore in the marine waters assessment these three factors include those that need to be measured and considered. Biological indicators including plankton, benthos and nekton can be used to assess macro changes in ecological balance, biological indicators are an easy guide to monitor the occurrence of contamination. In the presence of environmental pollution, the species diversity will decrease and the food chain becomes simpler, unless fertility occurs, the existence of Flora and marine fauna can be used as biological indicators of marine pollution can be observed from species diversity, and the rate of growth of the structure. The diversity of flora and fauna of high sea waters ecosystems indicates that the quality of water is good / not contaminated. But otherwise if the diversity is small, then the marine waters are polluted <sup>(14)</sup>

Coral reefs are marine organisms that can be used as bioindicators of damage levels of undersea ecosystems. Because of the importance of coral reefs in the ecological balance of the sea, the Indonesian government issued a coral reef rescue policy. the Indonesian government the <sup>6</sup> launched the program rescue coral reefs or better known as "Coral Reef Rehabilitation and Management Program "(COREMAP). Therefore, in every study on marine conservation, coral reefs and seaweed become bioindicator level of damage to marine waters. Damage to coral reefs and seaweed can be caused by unsustainable human development activities in managing and responding to the effects of industrial development. The development of fish landing harbor base can certainly affect the ecological balance of undersea flora and fauna. This is stated by Cooperation Center for Coastal and Marine Resource Studies <sup>(15)</sup> The method commonly used in the observation of

coral reefs and seaweed is the transect method. The use of transect method in describing the condition of the reefs corals are usually presented in the form of a community structure that consists of data: percentage of live coral cover, percent of coral cover dead, number of genera, number of species, number of colonies, size of colony, abundance, frequency of presence, growth form, coral reef diversity index. Sampling methods on coral reefs and sea grasses in transects would be better if supported by documentation through photo capture under the sea. Bioindicator of the marine ecosystem is a seaweed ecosystem. Seaweed is one of the biological resources found in coastal and marine areas. One indicator of the ecosystem is the seaweed ecosystem. Seaweed is one of the biological resources found in coastal and marine areas. These resources can usually be found in waters associated with the presence of coral reef ecosystems. Natural seaweed can usually live on sand and dead corals, likewise with plankton, benthos and nekton in the ocean can be used as bio-indicator of marine water damage level, its organism diversity indicates the level of environmental damage. Pollution levels can alter the structure of the ecosystem and reduce the number of species in a plankton, benthos and nekton communities, so that the diversity is reduced. Thus, the polluted marine ecosystem diversity index is always smaller than the natural marine ecosystem. Diversity in a marine waters are usually expressed in the number of species present in the site. The greater the number of species will be the greater the diversity. The relationship between the number of species and the number of individuals can be expressed in the form of diversity index. Organism diversity was analyzed by utilizing the diversity index formula Shannon Wiener, while the classification of the degree of pollution waters. Tropical coastal environments are biologically diverse environments which is good, but with the entry of pollutants causing marine ecosystems to become degraded by human activity that has the potential to cause extinction. Therefore, efforts should be made to preserve the organisms at sea.

Analysis of seawater quality in chemical, physical and biological components can provide an initial picture of the ecological status of waters sarrang, especially after the activity of fish landing bases, as well as analysis of plankton and benthos diversity indices as marine organisms susceptible to ecological changes due to contaminant entry at the Sarrang stage. With the reduction of planton and benthic diversity in marine waters Sarrang can be used as an indicator of the occurrence of pollution or ecological disturbance.

Therefore, research is needed to find out the initial picture condition of sarrang waters, especially in the waters around the fish landing base in sarrang village, rebang-Indonesia. The results of the study are expected to be the basis for the development and spatial plan of the Sarrang region including the planning of the Rebang district, especially with the government's plan to develop larger fish landing bases. So that local governments need a basis in determining strategic environmental assessment policies, environmental impact analysis and sustainable spatial and regional plans based on a good environment. Overview studies to find out the picture of an ecosystem is very important<sup>(17)</sup>

## MATERIAL AND METHOD

1. **Materials.** Material used in research are scaled ropes, Water bottle, Nansen Bottle Sampler, Plankton Net, glassware, titration equipment, 250 ml sample bottles, and 300 ml, pH meters, GPS, mercury thermometer, oven, label paper, secchi disk, turbidimeter, analytical balance, oven and AAS ( Atomic Absorption Spectrophotometer), Eckman Grab. Sedgwick-Rafter, Cover glass, Pipette drops, Book Identification, Microscope, dark sample bottle, Sedgwick-Rafter Counting Cell. Aquabidest, HNO<sub>3</sub>, Larutan lugol iodine, formalin, Asam asetat, Larutan lugol iodine, kalium iodida p.a, kalsium karbonat, Pb solution, 5 ml HNO<sub>3</sub> dilution solution, plankton sample bottle, Plankton-net, Aquades, Formalin 4%,

## 2. Method

The sampling site has temperatures between 28-29 ° C, with an air pressure of 764-768 mm Hg, with a moisture content of 74.5% - 78.9%. Wind speed 0.45 m / s - 1.36 m / s

### a) Sampling For Sea Water Parameters

The method used in determining the sampling location for the parameters of seawater and ground water quality is "Purposive Sampling".<sup>(18)</sup> Data on seawater quality in the waters surrounding Fish Landing Base Sarang is taken from the measurement of seawater quality on 21-24 October 2016 at 5 measurement stations. Measurement of seawater quality is done in the morning - noon time is at 08.00 - 12.00 is at optimum range conditions. Read more location and data of seawater quality as follows:

**Table 1.** Location of Sea Water Quality Measurement in Fish Landing Base Sarang Waters

No	Measurement Location	Coordinates	Depth
1	AL-1	110° 40' 18.57" E 6° 44' 12.27" S	2,8 m
2	AL-2	110° 40' 23.37" E 6° 44' 6.208" S	4,8 m
3	AL-3	110° 40' 10.314" E 6° 44' 4.334" S	3,0 m
4	AL-4	110° 40' 30.91" E 6° 44' 14.917" S	4,0 m
5	AL-5	110° 40' 29.996" E 6° 44' 2.215" S	6,8 m

(Primary Data, 2017)

The quality standard referred to is Minister of Environment Regulation no. 51/2004, concerning standard of quality standard of port designation. Sampling of seawater quality ie sample water is put into the sample bottle from each station, then it will be analyzed in Center for Industrial Pollution Prevention Technology Semarang with method of Atomic Absorption Spectrometer.<sup>(19)</sup>

### Sample extraction

The identification of air samples in quantities made under standard operating procedures for the method and air sample analysis. One hundred milliliters of air samples were transferred to a 500 mL separation funnel, added 5 mg of powdered NaCl and pH measured. A 50 mL n-hexane was added to the sample and shaken for 10 minutes. Samples are allowed to settle for 30 minutes to increase the mass of two

phases. The top organic layer is collected in a 300 mL Erlenmeyer flask, while the bottom layer moves to a 500 mL separation funnel. The extraction was repeated twice using 50 mL n-hexane. The organic phase is clean by adding 100 mL of pure air and shaken for 3 minutes. The sample is left fixed to increase two phases and then insert the bottom layer. The extraction is repeated twice using 100 mL of pure air. The extract was added with 5 mg of anhydrous sodium sulfate and concentrated to about 1 mL using a vacuum evaporator operating at 25 rpm and 35 ° C. Sediment samples were dried at room temperature in the laboratory, milled by mortar and sifted with 2 mm pore filter. Twenty-gram sediment samples were placed in a 250 ml flaked flask, supplemented with 40 mL acetone and shaken for 30 min. After the skin reaches, the extract is poured. Each crude extract was then dissolved in 10 mL of hexane and cleaned by passing the chromatographic column released by the anhydrous florasil and sodium sulfate. The net extract is concentrated in the rotary evaporator up to about 2 mL.<sup>(19)</sup>

The sampling methods for marine water quality were based on the Indonesian National Standard (SNI) No. 06-6989.57:2008 of The Methods of Surface Water Sampling. The analysis was used Atomic Absorption Spectrophotometry Method and while Total Suspended Solid (TSS) analysis was used gravimetric method according to Indonesian National Standard and APHA standard.<sup>(20,21)</sup>

## b) Sampling of Plankton and Nekton

Sampling of marine flora and fauna in the study area in the form of plankton, benthos, and nekton at the location of waters around Fish Landing Base Sarang Sampling done on 21-24 October 2017. With the coordinates as follows 6 ° 41'44.28 "S 111 ° 38'18.36 "T. Analysis of plankton, benthos and nekton by using Saprobity index analysis and Shannon Winner diversity index.<sup>(22)</sup>

Sampling of plankton and benthos is done at the same point. The phytoplankton and zooplankton sample were taken using plankton net with mesh size of 30-50 µm for phytoplankton and 0.2 mm for zooplankton. Then, the sampel were preserved with 4-5% formalin solution. The identification of plankton were used identification key and benthos sample were taken by grab sampler. The sediment that had been taken were shifted in water by 5 mesh sieve (254 mm). The filtered material then preserved by 10% formalin solution that had been added with coloring solution. The sample were identified by identifying key. The Nekton sampling process is to take advantage of GPS, boats, and catching equipment while the plankton sample is using plankton-nets.<sup>(23)</sup> Lee, *et al.* (1978) The water samples containing plankton and nekton are then taken to the Laboratory to identify their saprobity levels. The map below shows the location of seawater quality sampling and biological sampling, as follows :



Figure 1. Location of Seawater Quality Sampling-Groundwater and Sea Water Biota

**RESULTS AND DISCUSSION****1. Sea Water Quality Components**

Based on the results of seawater quality analysis are as follows :

**Table 2.** Quality of Sea Water in Fish Landing Base Sarang waters <sup>(19)</sup>

No	Parameter	Unit	Result Analysis					Quality standards
			AL-1	AL-2	AL-3	AL-4	AL-5	
<b>Physic</b>								
1	Brightness (TCU)	True Colour unit U Colour	0,75	1	1	1	1,5	>3
2	smell	-	No smell	No smell	No smell	No smell	No smell	No smell
4	Total suspended solid	mg/L	122	134	138	113	106	80
5	Trash	-	yes	Nihil	yes	Nihil	Nihil	Nihil
6	Temperature	OC	33	32	32	26	29	Natural
7	Oil residue	-	Nil	Nil	Nil	Nil	Nil	Nil
<b>Chemistry</b>								
1	pH	-	8,1	6,1	8,2	8,1	8,1	6,5 – 8,5
2	Salinity	‰	33,1	33,1	33,1	33,1	33,2	Natural
3	Ammonia total	mg/L	655	534	464	589	519	0,3
4	H <sub>2</sub> S	mg/L	0,57	0,57	0,57	0,74	0,57	0,03
5	Phenol	mg/L	0,22	0,02	0,01	0,01	0,01	-
6	Detergent	mg/L	0,84	0,97	0,62	0,82	0,99	1
7	Fat Oil	mg/L	211	226	271	210	303	5
<b>PARAMETERS- METAL SOLUTIONS</b>								
1	Cadmium (Cd)	mg/L	<0,01	<0,01	<0,01	<0,01	0,02	0,01
2	Copper (Cu)	mg/L	<0,04	<0,04	<0,04	<0,04	<0,04	0,05
3	Lead (Pb)	mg/L	0,13	0,14	0,15	0,09	0,12	0,05
4	Zinc (Zn)	mg/L	0,05	0,07	0,03	0,03	0,05	0,1
<b>PARAMETER MICROBIOLOGY</b>								
1	Coliform	MPN/100mL	11 x 10 <sup>3</sup>	7 x 10 <sup>3</sup>	14x 10 <sup>3</sup>	4 x 10 <sup>3</sup>	11 x 10 <sup>3</sup>	1000

Data Source: Primary Data of measurement results in October 2017

Quality Standard \*: Permenlh No. 51/2004, Seawater Quality Standards For Seaports <sup>(19)</sup>

AL1: location of measurement at distance 100 m north of Fish Landing Base Sarang

AL 2: location of measurement at distance 300m north of Fish Landing Base Sarang

AL3: location of measurement at distance 300m west of Fish Landing Base Sarang

AL4: location of measurement on the East Fish Landing Base Sarang

AL5: location of measurement at 500m distance north of Port of Sarang

The quality standard referred to is Minister of Environment Regulation no. 51/2004, is the seawater standard of Ports. Based on the measurement data of seawater quality at 5 (five) measurement points, that the quality of sea water in the waters around Fish Landing Base Sarang has been contaminated by organic contamination.

**1. Sea Water Parameters That Exceed Standard Standards**

The parameters of seawater quality exceeding the specified standard limits are Colliform range from 4 x 10<sup>3</sup> to 14 x 10<sup>3</sup> (MPN / 100 mL), the amount of ammonia (mg / L) range from 519 ppm - 655 ppm, Phenol between 0,02 ppm – 0,22 ppm, H<sub>2</sub>S range from 0,57 ppm - 0,74 ppm. colliform range from 4 x 10<sup>3</sup> to 14 x 10<sup>3</sup> (MPN / 100 mL) the allowable standard is 1000, it is intended that the waters are in the bad category. this indicates that the sanitation around the Sarrang fish landing

base is in poor condition. Sanitation due to organic waste from domestic activity is very bad, the waste processing industry at the fish landing bases has not been well implemented. By because it is very necessary for domestic waste results of fishing activities meet the established requirements. Research conducted by Karbasdehi in 2017 on Indicator bacteria community seawater and coastal sediment: the Persian Gulf as a case. Indicator bacteria community seawater and coastal sediment: the Persian Gulf as a case, indicating that the condition of coliform bacteria is quite tinggi with range 2326 (MPN / 100 mL) <sup>(33)</sup>. Waste treatment is needed, especially coliform, ammonia, phenol so that waste with this parameter go down so as to meet the quality standards of sea water Phenol is an organic compound, which is toxic, and easily soluble in water, making it easy to cause pollution in the water. Phenols can be toxic and if water exposed to phenol contamination will result in decreased water quality and disturbance to aquatic ecosystems. Based on Regulation Of Environment Minister)no. 51/2004, regarding sea water quality standards for ports refers to polluted waters, the standard limit of phenol concentration in marine waters is 0.002 ppm. Phenol contamination of marine waters in the Fish Landing Base Sarang caused by anthropogenic activity of largely organic nature degraded into final compounds, including phenol compounds. Phenol has a negative effect on the environment, especially aquatic ecosystems, water sources for population and public health around the location Fish landing base, because it is toxic and carcinogens. Phenol is also known to contaminate groundwater sources because it is easily soluble in water so that the pollution of coastal waters by phenol needs to receive intensive attention. <sup>(19)</sup>

Phenol (C<sub>6</sub>H<sub>6</sub>O) is an organic compound having hydroxyl groups which is attached to the benzene ring. This compound is often also called a hydroxy benzene or carbolic acid compound, phenate monohydroxybenzene, phenic acid, phenylic acid, phenyl hydroxide, oxybenzene, benzenol, monophenol, phenyl hydrate, phenyl alcohol, and phenol alcohol. <sup>(24)</sup> The level of phenol toxicity varies greatly depending on the number of atoms or molecules attached to its benzene chain. For chlorinated phenols, the more chlorine atoms binded to the benzene chain the more toxic. Chlorophenol may be more toxic to aquatic biota is accumulated and more persistent than simple phenols. Simple phenols such as phenol, cresol and xylenol are easily soluble in water and are more easily degraded. Phenol compounds are pollutants that are often found in the ocean waters. The source of pollutants in the ocean comes from crude oil spills due to the activities of fishermen, fuel spills of ships. The presence of phenol compounds in the ocean can endanger marine life because phenols are toxic. <sup>(25)</sup>

The ammonia source in the waters is the result of breakdown of organic nitrogen and inorganic nitrogen from Fish Landing Base activity, especially the activity of Sarang fish auction, which produces a lot of organic and inorganic waste which then degraded (amoniifikasi) by marine microorganism causing marine pollution. Canadian Council of Ministers of the Environment (1999) Ammonia can be decomposed on the sediments at the bottom of the sea floor. Ammonia in water can be lost through the evaporation process because the ammonia partial pressure in the solution increases with increasing pH.

The higher the pH of sea water, the higher the toxicity of ammonia. Because most of the NH-shaped, while the ammonia-shaped molecule (NH<sub>3</sub>) is more toxic than the ion (NH<sub>4</sub><sup>+</sup>). Ammonia in the form of molecules can penetrate the cell membrane parts faster than the NH<sub>4</sub><sup>+</sup> ions. Ammonia in the waters of Sarang already in the category high enough that is in the range 519 ppm - 655 ppm (standard permitted ammonia is 0.3 ppm).

Hydrogen sulfide (H<sub>2</sub>S) is derived from the activity of protein decomposition is organic matter. H<sub>2</sub>S is the result of degradation of proteins and organic matter by marine microorganisms that then decompose on sediments in the seabed. The result of seawater quality analysis in Fish Landing Base Sarang was detected H<sub>2</sub>S is 0,57 ppm from seawater quality standard of H<sub>2</sub>S which allowed only 0,03 ppm. <sup>(26)</sup>. Bacteria that can be used as a water pollution bioindicator include Coliform, Fecal Coliform, Salmonella and Streptococcus Fecal. The coliform bacteria found in waters of the Sarang Fish landing base with high concentrations is 4 x 10<sup>3</sup> to 14 x 10<sup>3</sup> (MPN / 100 mL), exceeding the approved standard is 1000 (MPN / 100 mL), this indicates that coastal seas have contaminated <sup>(27)</sup>

Pollution of the sea by coliform is possible because sanitation in this place is very bad because the organic material is directly discharged into the Based on the measurement results at 5 points of measurement at the location of the Sarang Fish Landing Base, it detected a high enough ammonia concentration at the sampling site 1 that is in the waters at 100 m north of Fish Landing Base Sarang, with a depth of 2.8 m and this point is the location which is closest to the location of the center of fish sales and population of fishermen settlements, while the lowest ammonia levels are in the location of AL3 in the waters at a distance of 300 m from West Port Sarang at a depth of 3 m. At AL-3 location because there is no building with sampling depth of 3 meter so it shows lower value of Total Coliform compared to other location. Based on the data of the analysis, the quality of seawater in Sarang Fish Landing Base based on physical, chemical and microbiological analysis (e coli and total coliform), the waters in the bad category with some parameters have exceeded the quality standards set based on Regulations Minister of Health of the Republic of Indonesia The groundwater quality support data is also measured to see the comparison of groundwater quality and sea water quality, groundwater quality data can be seen in Table 3. The groundwater quality in the Base Areas of Fish Nest is taken from groundwater quality measurement on 21-24 October 2016 at the well in the area of Hive Fishing Base, the well water around Fish Auction Sarang. From these measurements, it is known that the well water quality has been contaminated with the parameters shown: Total Dissolved Residues, Hardness, Ammonia and Microbiology. parameters (total Coliform and E Coli). above the clean water quality standard set by the Regulation of the Minister of Health of the Republic of Indonesia, <sup>(28)</sup>. This indicates that the well water is contaminated. Groundwater quality support data show poor groundwater quality as follows:

**Table 3.** Groundwater Quality at Activity Sites (Minister of Health of the Republic of Indonesia, 2017)

No	Parameter	Unit	Result of Analysis	Water Quality Standards
<b>A. PHYSICS</b>				
1	Smell	No Smell	No Smell	No Smell
2	Turbidity	NTU	1,48	25
3	Dissolved Residues (TDS)	mg/L	66	1.000
4	Color	TCU	*,17	50
5	Flavors	Not Feeling	Not Feeling	Not Feeling
6	Temperature	°C	31	± 3 °C Air temperature
<b>B. CHEMISTRY</b>				
1	Iron (Fe)	mg/L	0,13	1
2	Hardness (CaCO3)	mg/L	4.092	500
3	Chloride (Cl)	mg/L	36,7	-
4	Manganese (Mn)	mg/L	<0,07	0,5
5	pH	-	7,5	6,5 – 8,5
6	Zinc (Zn)	mg/L	0,03	15
7	Sulfat (SO4)	mg/L	159,8	400
8	Copper (Cu)	mg/L	<0,04	0,05
9	Ammonia (NH3-N)	mg/L	370	-
<b>Organic Chemistry</b>				
1	Total Chromium	mg/L	<0,04	-
2	Cadmium (Cd)	mg/L	<0,01	0,005
3	Nitrite as N	mg/L	0,08	-
4	Nitrate as N	mg/L	4,41	1
<b>C. MICROBIOLOGY</b>				
1	Total Coliform (MPN)	MPN/100mL	20 x 10 <sup>3</sup>	50
2	E.Coli	MPN/100mL	7 x 10 <sup>3</sup>	0

Data Source: Primary Data, Soil Water Quality in the area around **Fish Landing Base Sarang**

Quality Standard: Minister of Health of the Republic of Indonesia, 2017

**2. Biological Components**

**Sea Flora and Sea Fauna**

The conditions of marine flora and fauna studied in the study area are plankton, benthos, and nekton. In the study area was taken based on sampling of primary data in the location of waters around the fish landing base conducted on 21 -24 October 2017. 21 -24 October 2017.

Plankton and Benthos, The condition of Plankton and Benthos in the study area is based on the results of Saprobitas analysis and its diversity index as follows (17) (table 4):

**Table 4.** Analysis of Saprobitas - Fish Landing Base Sarang

	Station				
	S1	S2	S3	S4	S5
<b>Poly-saprobic Group (A)</b>					
Sum (n A)	0	0	0	0	0
<b>α- Meso-saprobic Group (B)</b>					
<i>Chaetoceros</i> sp		12			12
<i>Navicula</i> sp.	6	6	6	6	
<i>Nitzschia</i> sp.	12	90	36	42	96
<i>Vorticella</i> sp	24				
<i>Stephanodiscus</i> sp		24			18
sum (nB)	42	132	42	48	126
<b>β-Mesosaprobic Group (C)</b>					
<i>Tabellaria</i> sp	72	108	24	66	54
sum (nC)	72	108	24	66	54
<b>Oligosaprobic Group (D)</b>					
sum (nD)	0	0	0	0	0
<b>Group E (without A, B, C dan D)</b>					
<i>Balanus tintinabulum</i>	6				6
<i>Ceratum furca</i>				12	
<i>Ceriodaphnia</i> sp			18		
<i>Corycaeus ovalis</i>	24	18	24	24	42
<i>Euplotes</i> sp				6	
<i>Frontonia</i> sp	54				
<i>Macrosetella gracilis</i>	6	12	25	18	6
			2		
<i>Macrocyclops fuscus</i>	54	78		78	25
					8
Larva <i>Portunus pelagicus</i>	168	90	16	56	84
			2		
<i>Protococcus</i> sp		6			
<i>Richterella</i> sp	42	42	6		18
<i>Spirotaenia condensata</i>			6		
Jumlah (nE)	354	246	46	19	41
			8	4	4
$SI = \frac{1(nA) + 3(nD) + 1(nB) - 3(nA)}{1(nA) + 1(nB) + 1(nC) + 1(nD)}$	1,0	1,00	1,0	1,0	1,0
	0		0	0	0
$TSI = \frac{1(nA) + 3(nD) + 1(nB) - 3(nA)}{1(nA) + 1(nB) + 1(nC) + 1(nD)}$	0,2	0,49	0,1	0,3	0,3
$\times \frac{nA + nB + nC + nD + nE}{nA + nB + nC + nD}$	4		2	7	0

Description: (\*\*)= β - Mesosaprobic, Description Polluted mild to moderate (17,22)

Source: Analysis Results, 2017



Ecological diversity is expressed by index value, as statistical analysis of ecological diversity so as to estimate the abundance of a plankton and benthos in making a conclusion especially in comparing the divers<sup>46</sup> among some communities. Environmental changes have a strong effect on a single organism structure, including species richness and cor<sup>46</sup> sition. Pancton and benthic macroinvertebrates included in this biological component are very sensitive to change, because it includes various taxa with various cha<sup>46</sup> racteristics that are sensitive and tolerant species. Reducing the number of species along the environmental gradient has been linked to various environmental change factors, mainly due to the entry of pollutants due to human activities. Therefore, the diversity of planton and benthos can be used as an indicator of the quality of a marine waters through the calculation of the saprobites index.<sup>(29)</sup>

The result of saprobites analysis showed that the waters around Fish Landing Base Sarang were classified as mild to moderate. The sources of pollution include loading-unloading activities-fish auction in Fish Landing Base, and waste disposal by citizens to the sea. These findings support the results of water quality analysis, whereby several water quality parameters have been shown to exceed the threshold.

Pollution that occurs in the waters around the Fish Landing Base Sarang cause disruption to the biota, either against plankton, benthos or nekton. Several types of biota that can tolerate the pollution conditions will be able to survive, but the type of biota that is not able to tolerate pollution will experience a decrease in abundance due to death and migrate away from contaminated locations. Therefore, the management of the aquatic environment around Fish Landing Base Sarang is expected to improve the condition of waters and aquatic biota that have habitat in the waters around Fish Landing Base Sarang, either plankton, benthos or nekton.

The dominant types of plankton in the waters surrounding the project include *Macrocylops fuscus*, *Portunus pelagicus* (crab larvae), *Nitzchia sigma*, *Frontonia sp*, and *Tabellaria sp*. *Nitzchia sp* is one type of marine plankton that has a high tolerance to extreme environments, including having a high resistance in the aquatic environment that has contamination of organic matter. *Tabellaria sp* belongs to one group of oligosaprobic organisms as an indicator of the waters being at the level of mild or uncontaminated contamination. Saprobites analysis results show that the waters around Fish Landing Base Sarang have been contaminated with mild to moderate levels of pollution<sup>(30)</sup>

### Nekton

The results of the nekton survey indicate that the dominant fish species found in the waters around the project are peti fish (*Leiognathus equulus*) and kamojan fish (*Nemipterus nematophorus*). The diversity index is 1.74 (medium category), and the uniformity index is 0.837 (stable category).<sup>(30)</sup>

Makrozoobentos survey results before the project showed that benthos type encountered was dominated by *Anadara sp* type, followed by *Pleuricera sp* and *Unio sp*. *Anadara sp* is a tropical animal that lives in shallow waters to a depth of 20 m. *Anadara sp* including filter-feeder animals with main meals are detritus, as well as relatively resistant to pollution levels.

Nekton is a marine organism that can move or swim by itself in water so it does not depend on strong ocean currents or wind motion caused by wind. Nekton includes fish, squid, shrimp, crabs, mammals and marine reptiles. Location of nekton samples around Fish Landing Base Sarang, ie at coordinate points 6 ° 41'44.28 "S 111 ° 38'18.36" T. The tools used in this nekton sampling process are GPS, boats, and arad traps. Sampling is done at three different points.<sup>(31)</sup>

The degree of species diversity consists of two components: the number of species present (generally leading to species richness) and relative species abundance lead to uniformity. Diversity is generally measured using the distribution patterns of some abundance (individual or productivity) sizes between species. Data of biota number obtained at each point then analy<sup>67</sup> with index of diversity, and index of uniformity by using formula as follows.

$$H' = - \sum_{i=1}^n P_i \ln P_i$$

$$P_i = n_i/N$$

Where H' is the index of diversity, <sup>48</sup> is the number of individuals per species, and N is the total number of all species. The criteria for diversity index are as follows.

H' < 1 indicates a low level of species diversity

1 < H' < 3 indicates a moderate level of species diversity

H' > 3 shows a high level of species diversity

While the uniformity index can be calculated by the following formula :

$$E = H' / H_{max}$$

$$H_{max} = \ln S$$

Where E is the uniformity index and S is the total number of species. The value of uniformity index ranges from 0-1. If the value of E < 0.20 can be said unstable type uniform conditions, whereas if the value of E is 0.21 < E < 1 then it can be said stable type uniformity conditions.

Based on the sampling of nekton at three different points, there were 8 different fish species with 172 individuals total. Kamojan fish and petek are the most common fish species in the sampling

Table 5. Results Of Nekton Analysis









No	Name of Species	n <sub>i</sub>	P <sub>i</sub>	Ln P <sub>i</sub>	P <sub>i</sub> Ln P <sub>i</sub>
1	Buntek ( <i>Arothron hiudus</i> )	9	0,05232558	- 2,9502699	-0,15437459
2	Cumi cumi ( <i>Loligo sp</i> )	12	0,06976744	-2,66258783	-0,18576194
3	Rejung ( <i>Sillago sihama</i> )	8	0,04651163	-3,06805294	-0,14270014
4	Bawal Putih ( <i>Pampus argenteus</i> )	1	0,00581395	-5,14749448	-0,02992729
5	Badong ( <i>Eubleekeria jonesi</i> )	3	0,01744186	-4,04888219	-0,07062004
6	Petek ( <i>Leiognathus equulus</i> )	24	0,13953488	-1,96944065	-0,27480567
7	Kamojan ( <i>Nemipterus nematophorus</i> )	105	0,61046512	-0,49353413	-0,30128537

No	Name of Species	$n_i$	$P_i$	$\ln P_i$	$P_i \ln P_i$
8	Kakap Pungar ( <i>Lutjanus johni</i> )	10	0,05813953	-2,84490938	-0,16540171
	Sum	172	1	-23,1851715	-1,32487675
H' (index of diversity)		1,324877 (moderate)			
Hmax		2,079442			
E (uniformity index)		0,637131 (stable)			

66

Based on the results of the analysis indicated that the diversity index was 1.32 (moderat category) and the uniformity index was 0.63 (stable category) <sup>(17)</sup>. Description of fish species for <sup>(21)</sup> at the time of sampling by photo mapping method <sup>(32)</sup>. Its can be seen in the following table.

**Table 6.** Types of Identified Fish

No	Name	figure	Sum
1	Buntek ( <i>Arothron hiidus</i> )		9
2	Cumi cumi ( <i>Loligo sp</i> )		12
3	Rejung ( <i>Sillago sihama</i> )		8
4	Bawal Putih ( <i>Pampus argenteus</i> )		1
5	Badong ( <i>Eubleekeria jonesi</i> )		3
6	Petek ( <i>Leiognathus equulus</i> )		24
7	Kamojan ( <i>Nemipterus nematophorus</i> )		105
8	Kakap Pungar ( <i>Lutjanus johni</i> )		10

The results of this study are a general description of the marine ecological conditions of the latest sarrang and can be used as a baseline for other activities, especially environmental impact assessment studies and environmental feasibility studies to determine the predicted impacts that will arise if the fish landing ports will be expanded and expanded. Rembang district government as a stake holder area can utilize the results of this study for the planning of marine territorial waters district rembang, central java province

## CONCLUSION

Based on the analysis of marine water quality Sarang shows the waters in polluted condition, with The parameters of seawater quality beyond the prescribed standard limit are lead (Pb) ranging from 0.9 ppm - 0.15 ppm (mg / L), Colliform range from  $4 \times 10^3$  to  $14 \times 10^3$  (MPN / 100 mL), ammonia (mg / L) ranged from 519 ppm - 655 ppm, Phenol between 0.02 ppm - 0.22 ppm, H2S range from 0.57 ppm - 0.74 ppm, while the analysis Saprobik Index = 0.1, Tropic Saprobik Index = 0.24-0.49 and Nekton's Diversity Index is 1.74 shows the waters are also in polluted condition

## SIGNIFICANCE STATEMENTS

This study is a case study of recent studies on the ecological conditions of waters sarrang, especially its impact due to landing activity of fishing boats. As is known that the waters of Sarrang including in the conservation area Karimunjawa marine park that is protected by the Indonesian government. The results of this study provide the basis for the stakeholders who are authorized in the management and monitoring of ecological conditions of waters sarrang and as a frame of reference in the study of environmental impact analysis, especially in the field of marine waters assessment

## ACKNOWLEDGMENTS

This research is a part of research from Project analysis of environmental impact of development plan of Sarrang fish landing base center which is conducted between Mitra Adhi Pranata Company with Marine Development And Fisheries Of District Of Rembang no. 00022 / K-PA / 3.01.01.01 / 06/2017, Date 15 June 2017

The author is very grateful to many thanks to Mrs Poerna Sri Oetari as the President Director of PT Mitra Adhi Pranata who has allowed writers to conduct research at the Sarang and Mr. Aji as research assistant. We are very grateful to the head of department of Development And Fisheries Of District Of Rembang so that research can run well.

## REFERENCES

- [1] Cooperation Center for Coastal and Marine Resource Studies, 2000. Bogor Agricultural University and the Indonesian Institute of Sciences. Concept of Policy

- and Strategy National Management of Indonesian  
15]al Reefs. Bogor
- [2] Minister of Marine Affairs 15]d Fisheries of the Republic of Indonesia, 2014. Decree of the Minister of Marine Affairs and Fisheries of the Republic of Indonesia Number 45 / Kepmen-Kp / 2014 Concerning the Master Plan of National Fishery Port, Jakarta 65
- [3] Central Java Regulation of Governor, 2017. Central Java Governor Regulation No. 10 of 2017 Concerning Karimunjawa Port Master Plan. The Provincial Government of Central Java 2] Semarang,
- [4] Isworo and Oetari, 2017. Water Quality Status of River Donan due to Operational Refinery Pertamina Unit IV Cilacap-Central Java- Indonesia. Asian Journal of Biology 4(3): 1-10, 2017; Article 30 AJOB.36619 ISSN: 2456-7124
- [5] James W. Nybakken and Mark Bertness (2015), Marine biology : an ecological approach. 6th ed. San Francisco : Pearson/B 7] amin Cummings, 2005.
- [6] John R. Gray. 2015. Total Suspended Solids Data A Critical Evaluation. G. Douglas Glysson U.S. Geological Survey 412 National Center, Reston, Va 20192 Gglysson@Usgs.Gov John R. Gray U.S. Geological Survey 415 National Center, Reston, Va 20192 2
- [7] Minister Of The Environment, 2004. Decree Of The Minister Of Environmental State Minister Number 51 Year 2004 Regarding Quality Standard Of Sea State 23]hister For Environmental State. Jakarta 2004
- [8] Timothy M. Lenton and Andrew J. Watson. 2000. Regulation Of Nitrate, Phosphate, And Oxygen In The Ocean. Schoolo F Environmental Ciencesu, 51]ersity Of East Anglia,Norwich,United Kingdom Global Biogeochemic• Cycles, Vol. 14, No. 1, Pages 225-248, March 2000. 43
- [9] EPA, 2012. Appendix Environmental Protection Agency Response to Recommendations from Selected 39]C Reports: Policy, Activity, and Practice. Jakarta
- [10] Xiaodong Zhang, Lianbo Hu, and Ming-Xia He. 19]9. Scattering By Pure Seawater: Effect Of Salinity Department Of Earth System Science And Policy, University of North Dakota, Grand Forks, ND 58202, USA Ocean Remote Sensing Institute, Ocean 22]iversity of China, Qingdao, 266001, China
- [11] UNESCO, WHO and UNEP. 1996. Water Quality Assessments - a Guide to Use of Biota, Sediments and Water in Enviromental Monitoring. Second Edition. Edited by Chapman, Deborah. Publishing by F & FN Spon, London 25
- [12] Bo Barker Jorgensen and Friedhelm Bak. 1991; Pathways and Microbiology of Thiosulfate Transformations and Sulfate Reduction in a Marine Sediment (Kattegat, Denmark). Appl. Environ. 23]robiol, vol. 57 no. 3 847-856 27
- [13] Timothy M. Lenton and Andrew J., 2000. Regulation of nitrate, phosphate, and oxygen in the ocean. Global Biogeochemic Cycles, V 50]4, No. 1, Pages 225-248, March 2000 Watson School of Environmental Sciences, University of East Anglia, Norwich, United Kingdom
- [14] 2]woro\*, Poerna Sri Oetari , and N. A. Indah, 2017. Water Quality Status of River Donan due to Operational Refinery Pertamina Unit IV Cilacap-Central Java- Indonesia Asian Journal of Biology 4(3): 1-10, 2017; Article no.AJOB.36619 ISSN: 2456-7124. Indonesia 31
- [15] Herb Ward. 2015. Water Quality, Sediments, Sediment Contaminants, Oil and Gas Seeps, Coastal Habitats, Offshore Plankton and Benthos, and Shellfish. Rice University, H 38]on, TX, USA
- [16] Cadman, Cary Anne. 2017. Indonesia - Coral Reef Rehabilitation and Management Program- Coral Triangle Initiative (COREMAP-CTI) : P127813 - Implementation Status Results Report : Sequence 06 2]nglish). Washington, D.C. : World Bank Group
- [17] Kathleen A. Nolan and Jill E. Callahan. 2005. Beachcomber Biology: The Shannon- Weiner Species Diversity Index St. ABLE 2005 Proceedings Vol. 27 Francis College 180 Remsen St. Brooklyn, NY 11201 29
- [18] Andrea Crampton and Angela T. Ragusa 2016. Exploring Perceptions and Behaviors about Drinking Water in Australia and New Zealand: Hydrology 2016, 3, 8; doi:10.3390/hydrology 3010008 2
- [19] Decree of State Minister of Environment, 2004. Ministerial regulation Number 51 Year 2004 Concerning Water Quality Standard of State Minister of Environment Date: 8 April 2004. Jakarta
- 26
- [20] APHA. 1992. Standard methods for the examination of water and wastewater, 18<sup>th</sup> edition. American Public Health Association. Washington D.C. 2
- [21] Indonesian National Standard. 2017. SNI 06-6989.3-2004 Water and Waste Water- Part 3: Total Suspended Solids (TSS) Suspension Method Gravimetry.
- [22] Davide Tagliapietra and Marco Sigovini, 2012. Saprobity: A unified view of benthic succession models for coastal lagoons. Hydrobiologia 686(1) • May 2012 with 199. Italian National Research Council. 33
- [23] Goswami SC. 2004. Zooplankton methodology, collection & identification – A field manual. Nation Institue of Oceanography. Dona Paula, Goa 37
- [24] Fumihisa Kobayashi 24]eruyamaki And Yoshito Shinakamura, 2012. Biodegradation Of Phenol In Seawater Using Bacteria Isolated From The Intestinal Contents Of Marine Creatures International Biodeterioration & Biodegradation Volume 69, April 2012, Pages 113-118 9
- [25] J. Michałowicz and W. Duda 2007. Phenols – Sources and 28]icity. Polish J. of Environ. Stud. Vol. 16, No. 3 . University of Łódź, Faculty of Biology and Environment Protection, Banacha 12/16 str., 90-237 Łódź, Poland

- [26] Urszula Janas, and Anna Szaniawska, 1995. The influence of hydrogen sulphide on macrofaunal biodiversity in the Gulf of Gdańsk\* *Oceanologia*, No. 38 (1) pp. 127–142, 1996. PL ISSN 0078–3234 Institute of Oceanography, Gdańsk University, Gdynia
- [27] Partha Pal 2014. Detection of coliforms in drinking water and its effect on human health -International Letters of Natural Sciences Online: 2014-06-30 ISSN: 2300-9675, Vol. 17, pp 122-131 doi:10.18052/www.scipress.com/ILNS.17.122 © 2014 SciPress Ltd., Switzerland
- [28] Minister of Health of the Republic of Indonesia, 2017. Regulation of the Minister of Health of the Republic of Indonesia Number 32 Year 2017 on Standards of Quality Standard of Environmental Health and Water Health Requirements for Sanitation Hygiene, Swimming Pool, Solus Per Aqua, and Public Bath
- [29] Lorenzo Fattorini 2009. *Environmetrics - Statistical Analysis Of Ecological Diversity - Dipartimento Di Metodi Quantitativi*, Università Di Siena, Italy
- [30] Onyema IC. 2013. Phytoplankton bio-indicators of water quality situations in the Iyagbe Lagoon, South-Western Nigeria. Department of Marine Sciences, University of Lagos, Akoka, Lagos, Nigeria; 2013
- [31] Eduardo Ríos-Jara, Ceciel-M. Navarro-Caravantes, Cristian-M. Galván-Villa, and Ernesto Lopez-Urriarte, 2009. Bivalves and Gastropods of the Gulf of Tehuantepec, Mexico: A Checklist of Species with Notes on Their Habitat and Local Distribution. *Hindawi Publishing Corporation Journal of Marine Biology* Volume 2009, Article ID 176801, 12 pages doi:10.1155/2009/176801
- [32] Bennett and T.C. Lantz, 2014. Participatory photomapping: a method for documenting, contextualizing, and sharing indigenous observations of environmental conditions \* School of Environmental Studies, University of Victoria, PO Box 3060 STN CSC, Victoria, British Columbia V8W 3R4, Canada
- [33] Karbasdehi et al. 2017. Indicator bacteria community in seawater and coastal sediment: the Persian Gulf as a case. *Journal of Environmental Health Science & Engineering* (2017) 15:6 DOI 10.1186/s40201-017-0266-2

# Overview the Condition of Seawaters Around the Fish Landing Base in Sarang Indonesia

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

12%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Shahida Begum, Reshma Banu, Ali Ahamed, B. D. Parameshachari. "A comparative study on improving the performance of solar power plants through IOT and predictive data analytics", 2016 International Conference on Electrical, Electronics, Communication, Computer and Optimization Techniques (ICEECCOT), 2016  
Publication 2%
- 2 [www.sdiarticle2.org](http://www.sdiarticle2.org)  
Internet Source 2%
- 3 R. Maru, I. I. Baharuddin, N. Badwi, S. Nyompa, Sudarso. "Analysis of Water Well Quality Drilling Around Waste Disposal Site in Makassar City Indonesia", Journal of Physics: Conference Series, 2018  
Publication 1%
- 4 [shodhganga.inflibnet.ac.in](http://shodhganga.inflibnet.ac.in)  
Internet Source 1%
- 5 [pcbassam.org](http://pcbassam.org)

Internet Source

<1%

6

[docplayer.net](http://docplayer.net)

Internet Source

<1%

7

[vwrrc.vt.edu](http://vwrrc.vt.edu)

Internet Source

<1%

8

[ijcrbp.com](http://ijcrbp.com)

Internet Source

<1%

9

Submitted to Universiti Teknologi MARA

Student Paper

<1%

10

[emrlibrary.gov.yk.ca](http://emrlibrary.gov.yk.ca)

Internet Source

<1%

11

[healthyfood-list.blogspot.com](http://healthyfood-list.blogspot.com)

Internet Source

<1%

12

T.D. Bennett, T.C. Lantz. "Participatory photomapping: a method for documenting, contextualizing, and sharing indigenous observations of environmental conditions", *Polar Geography*, 2014

Publication

<1%

13

P YILDIRIM. "MONITORING AND EVALUATION OF COASTAL WATER QUALITY PARAMETERS IN FETHIYE BAY, TURKEY", *Applied Ecology and Environmental Research*, 2019

<1%

14

[www.sciencedomain.org](http://www.sciencedomain.org)

Internet Source

<1%

---

15

[eprints.unimal.ac.id](http://eprints.unimal.ac.id)

Internet Source

<1%

---

16

Davide Tagliapietra, Marco Sigovini, Paolo Magni. "Saprobity: a unified view of benthic succession models for coastal lagoons", *Hydrobiologia*, 2012

Publication

<1%

---

17

Arif Susanto, Purwanto Purwanto, Edi K. Putro, Fanny Yuliasari. "Drinking Water Risk Management Plan: A Case Study in the Ore Processing Industry", *Safety*, 2019

Publication

<1%

---

18

Nani Radias Tuti, Dwi Ningsih Susilawati, Nur Hasni. "The ENDOPHYTIC FUNGI COMMUNITIES FROM *Centella asiatica* OF BENGKULU ACCESSION", *Fungal Territory*, 2019

Publication

<1%

---

19

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1%

---

20

[www.actasatech.com](http://www.actasatech.com)

Internet Source

<1%

---

[www.scribd.com](http://www.scribd.com)

21	Internet Source	<1%
22	Submitted to University of Venda Student Paper	<1%
23	Submitted to Heriot-Watt University Student Paper	<1%
24	pub2.db.tokushima-u.ac.jp Internet Source	<1%
25	www.koreascience.or.kr Internet Source	<1%
26	www.ecy.wa.gov Internet Source	<1%
27	Submitted to UC, Irvine Student Paper	<1%
28	www.pjoes.com Internet Source	<1%
29	Submitted to Australian Catholic University Student Paper	<1%
30	documents.mx Internet Source	<1%
31	www.springer.com Internet Source	<1%
32	enviscecb.org Internet Source	<1%



---

33	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1%
34	<a href="http://www.docstoc.com">www.docstoc.com</a> Internet Source	<1%
35	D M Nuriasih, S Anggoro, Haeruddin. "Saprobic analysis to Marina coastal, Semarang city", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1%
36	A Tjahjono, O Wahyuni, S Purwantini. "The Assessment of Biological and Pollution Index of Estuaries Around Port of Tanjung Emas Semarang", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1%
37	<a href="http://www.elementascience.org">www.elementascience.org</a> Internet Source	<1%
38	<a href="http://yz-car.ci.cqvip.com">yz-car.ci.cqvip.com</a> Internet Source	<1%
39	<a href="http://aoip.osa.org">aoip.osa.org</a> Internet Source	<1%
40	Submitted to Asian Institute of Technology Student Paper	<1%
41	<a href="http://jehse.biomedcentral.com">jehse.biomedcentral.com</a> Internet Source	<1%

---

---

42 Karolina Pierścieniak, Justyna Grzymała, Maciej Wołowicz. "Differences in reproduction and condition of *Macoma balthica* and *Mytilus trossulus* in the Gulf of Gdańsk (Southern Baltic Sea) under anthropogenic influences", *Oceanological and Hydrobiological Studies*, 2010  
Publication

---

43 [www.nap.edu](http://www.nap.edu)  
Internet Source

---

44 [www.isca.in](http://www.isca.in)  
Internet Source

---

45 [bmcmicrobiol.biomedcentral.com](http://bmcmicrobiol.biomedcentral.com)  
Internet Source

---

46 [journal.ipb.ac.id](http://journal.ipb.ac.id)  
Internet Source

---

47 [shareme-share.blogspot.com](http://shareme-share.blogspot.com)  
Internet Source

---

48 Submitted to The University of the South Pacific  
Student Paper

---

49 Eduardo Ríos-Jara, Ceciel-M. Navarro-Caravantes, Cristian-M. Galván-Villa, Ernesto Lopez-Uriarte. "Bivalves and Gastropods of the Gulf of Tehuantepec, Mexico: A Checklist of Species with Notes on Their Habitat and Local

## Distribution", Journal of Marine Biology, 2009

Publication

---

50

[www.esig.ucar.edu](http://www.esig.ucar.edu)

Internet Source

<1%

---

51

Timothy M. Lenton. "Redfield Revisited 1. Regulation of Nitrate, Phosphate, and Oxygen in the Ocean", Global Biogeochemical Cycles, 2000

Publication

<1%

---

52

Ashari Rachmat, Budianto Emil, Herdiansyah Herdis, Karuniasa Mahawan, A Saragi Putri. "Minimize the impact of waste pollution in ship repair processes to Improving Shipyard Industrial Infrastructure Sustainability", E3S Web of Conferences, 2018

Publication

<1%

---

53

Submitted to Program Pascasarjana Universitas Negeri Yogyakarta

Student Paper

<1%

---

54

Submitted to Federal University of Technology

Student Paper

<1%

---

55

Ganeshkumar Arumugam, Rajaram Rajendran, Vinothkumar Shanmugam, Rameshkumar Sethu, Mathivanan Krishnamurthi. "Flow of toxic metals in food-web components of tropical mangrove ecosystem, Southern India", Human and Ecological Risk Assessment: An

<1%

56

[www.oalib.com](http://www.oalib.com)

Internet Source

<1%

57

Submitted to University of Newcastle

Student Paper

<1%

58

M P Wardani, A Fahrudin, F Yulianda. "Analysis of Successful Strategy to Develop Sustainable Marine Ecotourism in Gili Bawean Island, Gresik, East Java", IOP Conference Series: Earth and Environmental Science, 2017

Publication

<1%

59

[patents.google.com](http://patents.google.com)

Internet Source

<1%

60

P. Cady, T. B. Boving, B. S. Choudri, A. Cording, K. Patil, V. Reddy. "Attenuation of Bacteria at a Riverbank Filtration Site in Rural India", Water Environment Research, 2013

Publication

<1%

61

Submitted to University of Warwick

Student Paper

<1%

62

G Chen. "Utilization of oxygen in a sanitary gravity sewer", Water Research, 2000

Publication

<1%

63

Submitted to Ohio University

Student Paper

<1%

---

64

Submitted to Cranfield University

Student Paper

<1%

---

65

Submitted to Yeungnam University

Student Paper

<1%

---

66

[www.ijsrp.org](http://www.ijsrp.org)

Internet Source

<1%

---

67

[www.zoologie.upol.cz](http://www.zoologie.upol.cz)

Internet Source

<1%

---

68

[meetings.aps.org](http://meetings.aps.org)

Internet Source

<1%

---

69

[media.neliti.com](http://media.neliti.com)

Internet Source

<1%

---

70

Sunoko, Rahmadi, and Hsiang-Wen Huang.  
"Indonesia tuna fisheries development and  
future strategy", Marine Policy, 2013.

Publication

<1%

---

71

Submitted to Sumner High School

Student Paper

<1%

---

---

Exclude quotes      Off

Exclude matches      < 8 words

Exclude bibliography      Off

# Overview the Condition of Seawaters Around the Fish Landing Base in Sarang Indonesia

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---